

Sample description

Table 1 – analyzed sample description.

Description	Active ingredient	Nominal value
NAND 300 300 mg/ml 10 ml UFC Pharm (pic 1)	Nandrolonedecanoate	300 mg/ml



Pic. 1 – Sample appearance.

Sample preparation for analysis

Table 2 – Sample preparation.

Preparation
50 ul of the sample was mixed with 1500 ul dichloromethane in a vial.

Analysis

Chromatographic separation (GC).

Column: Tr-5ms, 30 m.

Carrier gas: helium.

Flow rate: 1.2 ml/min.

Split flow rate: 12 ml/min.

Split: 10.

Injector: PTVinjector, final temperature (transfer) 290 °C.

Inject 1ul.

Split program in table 3.

Table3 – splitprogram.

Ramp	Rate (°C/min)	Temp, °C	Hold
Initial		35	1.70
Ramp 01	30	41	2.00
Ramp 02	20	310	15


Mass-spectrometricdetector (ionitrap) wasusedfordetection (m/z 50-650).

Identification: DBNist, librarySWDrug.

.

Sample.

RT: 0.00 - 31.01



NL:
5.23E6
TIC MS
sample_13

Pic 2 – Chromatogram of the sample. Retention time of active ingredient is 20.82 мин.

Table 4 – Identified ingredients of the sample.

Retention time	Substance	Peak area	Response factor	content, mg/ml
13.10	Ethylolate	8 692 863	1.4	
20.82	Nandrolonedecanoate	5 026 438		299

Results

Table 5 – Analysis results.

Active ingredient	Content of an active ingredientmg/ml	Foundexipients
Nandrolonedecanoate	299	Ethyl oleate.